

# Exercise Prescription for Older Adults With Osteoarthritis Pain: Consensus Practice Recommendations

*A Supplement to the AGS Clinical Practice Guidelines on the Management of Chronic Pain in Older Adults*

American Geriatrics Society Panel on Exercise and Osteoarthritis

## INTRODUCTION

In response to mounting evidence that a program of increased physical activity is a useful component in the management of osteoarthritis (OA) in older adults, the Board of Directors of the American Geriatrics Society (AGS), along with a multidisciplinary panel of experts, recommended that the AGS take the lead in promoting exercise prescription for OA patients in the primary care setting. This project, culminating in the publication of these practice recommendations, parallels the Society's clinical practice guidelines on the management of chronic pain in older persons.<sup>1</sup>

The purpose of this document is to provide an evidenced-based review that explains why a physically active life style benefits older adults with OA and to provide practical strategies and exercise guidelines for this expanding patient population. These practice recommendations are derived from the existing literature and by consensus among a panel of experts from many disciplines: geriatrics, internal medicine, orthopedics, physical therapy and rehabilitation, exercise physiology, nursing, and pharmacy. A literature search involving a full-text computer search of *Index Medicus* and MEDLINE using the terms *osteoarthritis*, *exercise*, and *aging* was first conducted. An extensive manual search using the bibliographies of the publications located through the computer search was also undertaken. A study was included in this review if the publication made an implicit or explicit claim regarding osteoarthritis or research designed to evaluate the effects of exercise on physiologic or functional parameters in older adults. Members of the multidisciplinary panel reviewed successive drafts of the report summarizing their findings, and the final draft was submitted for review and comment by experts routinely involved in the care of older adults.

Americans 65 years or older represent an expanding proportion of the United States (U.S.) population, and their numbers will increase rapidly as the baby-boom generation ages.<sup>2,3</sup> Although the majority of older people in

the U.S. are healthy and physically active, others suffer with chronic illnesses and require some assistance (family, friends, and public support systems) to manage their everyday lives.<sup>4,5</sup> Approximately one quarter of all patients seen by primary care physicians present with musculoskeletal conditions<sup>6</sup> and, among those age 65 years and older, the most prevalent articular disease is OA.<sup>7</sup> Addressing the health care needs of this rapidly expanding population is a national priority.

The conspicuous presence of OA in the older population has many believing that chronic pain and functional difficulties are immutable consequences of aging. OA, the most common form of arthritis, is associated with considerable disability.<sup>8</sup> Symptomatic OA causes pain, limits daily activities, and reduces quality of life.<sup>8,9</sup> The majority of those burdened with OA are elderly; in fact, about half of all persons age 65 and over are affected by OA.<sup>10</sup> The fallacy that undercuts the mistaken belief that symptomatic OA is caused by aging is revealed in the following anecdote. An older man visits his doctor complaining of difficulty with getting out of a chair and walking because of persistent pain in one of his knees. The doctor replies, "Well you're 75, this is just part of growing old." The astute patient replies, "My other knee is just as old and it doesn't hurt."<sup>11</sup>

An emerging body of evidence shows that light- to moderate-intensity physical activity may play a preventive and possibly a restorative role in combating declines in health and functional capacity caused by chronic diseases such as OA.<sup>12-17</sup> Regular physical activity modifies risk factors for chronic diseases prevalent in the older population,<sup>18,19</sup> improves psychologic health,<sup>10</sup> and promotes functional independence.<sup>20-33</sup> Physical inactivity is recognized as a risk factor for many diseases prevalent in the older population (coronary artery disease, diabetes mellitus, and obesity), and increasing physical activity in sedentary OA patients may reduce morbidity and mortality.<sup>34-38</sup> Evidence indicates that quadriceps muscle weakness is a risk factor for knee OA, which is often the consequence of inactivity.<sup>35</sup> Furthermore, inactivity may contribute to the morbidity associated with a variety of other chronic diseases, most notably diabetes mellitus, cerebrovascular disease, coronary artery disease, congestive heart failure, osteoporosis, and depression. The superimposition of any of these

---

Reviewed and approved by the AGS Clinical Practice Committee and the AGS Board of Directors, May 2001.

Address correspondence and reprint requests to Nancy Lundebjerg, Senior Director, Professional Education and Publications, American Geriatrics Society, 350 Fifth Avenue, New York, NY 10118.

conditions on several age-related changes only compounds the negative effects of OA on the older person's functional independence. Encouraging regular exercise may reduce the physical impairments and the burden of comorbidities, and thus improve the OA patient's quality of life.<sup>36</sup>

Comprehensive management of the patient with OA should involve non-pharmacological interventions in combination with medications. Medications such as analgesics and nonsteroidal anti-inflammatory drugs (NSAIDs) should not be used alone as the primary therapy but instead should be used in conjunction with non-pharmacologic measures.<sup>11,39,40</sup> These include education about joint protection, weight-loss counseling for obese people, development of pain-coping skills, enhancement of social support, application of heat or cold to painful joints, exercises that strengthen muscles, and the use of a cane or a walker. Developing an exercise program aimed at alleviating pain and improving overall physical fitness is especially important, because the primary concern for many OA patients is maintenance of functional independence.<sup>12,41,42</sup>

### OA RISK FACTORS, INCIDENCE, AND PREVALENCE: IMPLICATIONS FOR TREATMENT

Osteoarthritis is a major cause of chronic pain and disability in the older population. Even though there is much that we still do not understand about the pathophysiology of OA, our current understanding is sufficient to direct therapeutic interventions. Research in the underlying mechanisms of OA have identified several risk factors. The data suggest that, as with atherosclerotic heart disease, the risk of developing symptomatic OA is influenced by the presence of multiple risk factors. Reducing or eliminating these risk factors may reduce the symptoms and disability associated with OA. Table 1 lists the major risk factors of knee OA in older adults.<sup>11</sup> Some factors, such as age, gender, and inheritance are immutable, but others are modifiable. Obesity, muscle weakness, joint laxity, and altered biomechanics are some risk factors potentially amenable to non-pharmacologic measures. An understanding of the ways these risk factors affect the course of OA provides clinicians with the rationale for targeting their interventions for OA patients and increases the likelihood that these patients will improve.

The incidence and prevalence of OA at different anatomic regions vary, depending on whether this condition is defined by clinical symptoms, radiologic findings, or a combination of the two.<sup>43-46</sup> Although all peripheral joints may be affected, OA of the knee has been the focus of many epidemiologic studies.<sup>9,47-55</sup>

Age is the most consistent risk factor for both radiographic and symptomatic OA at all articular sites.<sup>46</sup> The prevalence of OA increases after the age of 40 in women

and 50 in men.<sup>55,56</sup> OA affects about 50% of persons age 65 and older, and this prevalence increases to 85% in the group age 75 and older.<sup>57,58</sup> In the Framingham Osteoarthritis Study, Felson and colleagues<sup>56</sup> found that 27% of the people age 63 to 70 years had knee OA diagnosed radiographically, and among those 80 years or older, the prevalence increased to 44%. In 1997, researchers found that nearly 12% of people age 65 years and older said that their activities were limited because of arthritis.<sup>1</sup> This number is likely to grow proportionally as elderly people comprise an increasingly greater share of the U.S. population in the decades to come.

Gender also influences the prevalence and incidence of OA. Isolated hand and knee OA are common in women, whereas the prevalence of hip disease is higher in men.<sup>44,45,49,50,54</sup> Prospective, longitudinal studies have examined the relationship between body weight and OA. Data from the Framingham Knee Osteoarthritis Study, which followed 1,420 persons for more than 30 years, indicate that overweight men and women are at higher risk for developing symptomatic and radiographic OA than those less obese.<sup>53,59</sup> Similarly, both the Baltimore Longitudinal Study of Aging and the Swedish study demonstrated that obesity increased the risk for developing OA.<sup>60,61</sup> Felson and colleagues also reported that weight reduction reduces pain, further supporting the relationship between obesity and OA.<sup>62</sup> Although the exact mechanisms remain unclear, several investigators speculate that excessive body weight increases the biomechanical stresses across weight-bearing joints, which eventually results in cartilage damage.<sup>63-65</sup> Although some data support this teleologic hypothesis, a direct relationship between weight loss and reduced OA morbidity is less convincing. A small limited number of randomized clinical trials provide preliminary data suggesting that a reduction in OA symptoms is correlated more strongly with reduced body fat mass than with reduced total body weight.<sup>56,66</sup> Slemenda and colleagues<sup>67,68</sup> reported that reduced strength, relative to body weight, may play a role in the development of OA. These preliminary data could indicate that reduced total body fat and increased muscle strength may be relevant to the development of OA. These data suggest that interventions designed to strengthen the muscles and reduce total body fat may be effective methods for reducing pain and improving function in patients suffering with OA.

A history of joint trauma, the presence of bony deformities, or joint instability are also risk factors for OA.<sup>9</sup> Evidence from a variety of cross-sectional and longitudinal studies suggests that major trauma to a joint increases the risk for developing OA.<sup>69-73</sup> Consistently, the evidence suggests a strong relationship between joint damage and the development of OA later in life.

**Table 1. Major Risk Factors for Osteoarthritis**

Modifiable	Potentially Modifiable	Immutable
Obesity	Trauma	Age
Muscle weakness	Reduced Proprioception	Fender female > male
Heavy physical activity	Poor joint biomechanics (i.e., joint laxity)	Inheritance
Inactivity		Congenital (i.e., malformations)

Recent studies report that muscle weakness and reduced joint proprioception are risk factors for developing OA. In patients with knee OA, quadriceps weakness is common. Because of decreased joint stability and shock-absorbing capacity, muscle weakness contributes to disability.<sup>74</sup> Until recently, this disuse muscle atrophy was thought to develop because patients avoid loading painful joints. Slemenda and colleagues<sup>67</sup> examined this relationship between muscle strength and knee OA in a population of randomly selected community-dwelling older adults age 65 years and older. They reported that in those without a history of knee pain, isolated quadriceps weakness was strongly associated with radiographic knee OA. The findings from subsequent studies<sup>68,75,76</sup> suggest that quadriceps weakness may be a risk factor for knee OA.

Although proprioception declines with age, several reports demonstrate that diminished position sense contributes to the development of OA.<sup>77–80</sup> Pai and colleagues showed that knee proprioception was significantly diminished in older adults with knee OA when compared with their counterparts without OA.<sup>77</sup> Whether reduced proprioception causes or is a consequence of OA remains unknown. Highlighting the importance of these factors are the findings from several studies that demonstrated strengthening and aerobic exercise programs designed to improve muscle strength and joint proprioception reduce pain and improve mobility in patients with OA.<sup>75,76,81–83</sup>

The relationship between levels of physical activity and the risk of OA has been well studied. In general, moderate amounts of recreational physical activity do not increase the risk of OA.<sup>84–86</sup> However, participation in occupations requiring strenuous physical activity or intense competitive sports activity throughout life may contribute to the development of OA.<sup>59,87–89</sup> Nevertheless, the results of many studies suggest that older adults, even those with OA, can reduce their morbidity by regularly participating in moderate physical activity.

### **BENEFITS OF PHYSICAL ACTIVITY FOR OA PATIENTS**

Published reviews outline the effects of exercise training in the OA population.<sup>90,91</sup> The details of selected randomized, controlled studies are presented in Table 2,<sup>42</sup> which also lists many of the benefits of increased physical activity for OA patients. Although many researchers who have studied this question conclude that exercise training does not impact the pathological process of arthritis,<sup>30–32,60,76,82,92–95</sup> a notable and consistent finding across the OA literature is that exercise training does not exacerbate pain or disease progression and is effective in decreasing pain and improving function. Moreover, the evidence from well-controlled clinical trials suggests that regular physical activity can provide older OA patients with the same physical, psychologic, and functional benefits observed in the general population. Chief among the functional benefits produced by increasing physical activity is improved postural and gait stability, which may reduce falls in this at-risk population.<sup>76,96–100</sup> These findings are significant because emerging research data implicate muscle weakness as a risk factor for OA<sup>67,68,101</sup> and suggest that physical inactivity exacerbates disability in OA patients.<sup>102</sup>

Short-term studies show that aerobic exercise<sup>82,103</sup> and strength-training programs<sup>17,83,100</sup> effectively improve impor-

tant physiologic parameters related to functional capacity in older adults with OA. More recent randomized, controlled long-term trials confirm these earlier findings.<sup>75,74,102,104</sup> Given the positive health and functional benefits associated with exercise and the fact that inactivity contributes to disability,<sup>102</sup> it is evident that promoting physical activity should be an integral component of the management of OA.

### **MANAGEMENT OF SYMPTOMATIC OA**

To date, no definitive treatment or cure for OA has been identified. The management of OA includes patient education, therapeutic modalities, exercise, and medications in parallel. Treatment goals include pain control, maximizing functional independence and improving quality of life within the constraints imposed by both OA and comorbidities.<sup>40,105</sup>

#### **Patient Education and OA**

Patient education is an important component of effective arthritis rehabilitation. Counseling programs have been found to be effective in reducing the pain and disability associated with OA.<sup>106–110</sup> Formal community-based programs to which the primary practitioner can refer OA patients are also available in many locations. In addition, the Arthritis Foundation publishes educational brochures and videotapes for patients and, in many communities, offers courses that teach practical techniques to reduce pain and improve function and general health. The Arthritis Foundation maintains a Web site (<http://www.arthritis.org>) and can be reached at 1-800-283-7800.

#### **Therapeutic Modalities**

Modalities, such as heat, cold, sound, and electricity are adjunctive interventions that are used with exercise and medications. While little scientific data demonstrate the efficacy of any specific modality in OA treatment, topical applications of heat and cold have been used for thousands of years. The physiologic effects of thermal modalities include muscle relaxation and decreased pain. However, because adverse effects can occur with the application of heat and cold, a comprehensive medical evaluation is necessary prior to using these modalities. The physiologic effects and general precautions of these modalities are outlined in Table 3.

### **EXERCISE ASSESSMENT AND PRESCRIPTION**

Pain, swelling, limited range of motion, muscle weakness, postural or gait instability, and poor cardiovascular fitness are significant physical impairments associated with OA. Interestingly, sedentary people without arthritis have many of these same problems, which suggests that physical inactivity plays an important role in the symptoms and signs associated with OA. A potential barrier to recommending regular physical activity to patients with OA is the belief that exercise will exacerbate joint symptoms. The results of randomized, controlled clinical trials, however, indicate that increased physical activity does not produce or exacerbate joint symptoms and, in fact, confers significant health benefits.<sup>30,32,52,54,59,60,111</sup>

#### **Patient Screening**

A comprehensive evaluation is the initial step in designing a physical activity program individualized for the patient with OA. The information obtained provides the founda-

**Table 2. Effects of Exercise Training in Osteoarthritis Patients: Randomized, Controlled Trials**

Author, Year	Exercises (Modes, Volume, Frequency, Intensity)	Sample Population (Groups/Subjects [n], Age [yr], Disease Classification)	Outcome Measures (Physiological, Psychological, Functional, Disease Activity)	Results
Minor, 1989 <sup>82</sup>	Aerobic walking and aquatic exercise 12 wk, 3/wk, 30-min sessions, HR <sub>MAX</sub> 60%-80%	Walking, aquatic/n = 49 Control/n = 19 Age: 63.8 ± 8.6 Age range: 36-83	Physio: VO <sub>2</sub> max trunk flexibility FXN: 50-ft walk test Psych—AIMS: phys activity, pain anxiety, depression Dz: swollen joints, morning stiffness	↑ In exercisers ( $P \leq 0.01$ ) ↑ In aerobic exercisers ( $P \leq 0.01$ ) ↑ In aerobic exercisers ≈ 12% ( $P \leq 0.01$ ) ↓ In aerobic group ( $P \leq 0.01$ ) ↓ in aerobic group ( $P \leq 0.05$ ) ↓ No. swollen joints in aerobic group ( $P \leq 0.01$ )
Kovar, 1992 <sup>83</sup>	Supervised walking program 8 wk, 3/wk, 30-min sessions, intensity: NR	Walkers/n = 47 Age: 70.4 ± 9.1 Control/n = 45 Age: 68.5 ± 11.3	Fxn: 6-min walk Psych—AIMS: phys act'y disability pain medication	↑ Distance 70 m in exercise group but 17 m in control group ( $P \leq 0.001$ ) ↓ in exercisers ( $P < 0.001$ ) ↓ in exercisers ( $P < 0.003$ ) Slight ↓ in exercisers ( $P = 0.08$ ) Significant ↑ both exercise groups in comparison with controls Significant ↑ aerobic exercisers in comparison with controls
Ettinger, 1997 <sup>77</sup>	Aerobic training: 3-mo supervised walking then 15-mo home walking, 3/wk, 1-hr sessions; HF <sub>reserve</sub> : 50%-70% Resistance training: 3-mo supervised the 15-mo home program, 9 exercises of UE and LE muscle groups, 2 sets of 12 repeats 3 d/wk; intensity: NR Control: health education program	Aerobic training/n = 144 Age: 69 ± 6 Resistance training/n = 146 Age: 68 ± 6	Physio: Isokinetic strength of knee flexion at 30°/sec V <sub>o2</sub> max Psych: self-report of phys disability	Significant ↑ aerobic exercisers in comparison with controls Both exercise groups reported less disability than controls Both exercise groups did better than controls No difference in severity of OA among 3 groups Both exercise groups reported significantly less pain than controls Significant ↑ (20%) extension strength in exercise groups least affected knee at 30°/sec at 3 mos ( $P < 0.05$ ) ↑ (21%) extension strength in exercise groups least affected knee at 3 mos ( $P = 0.06$ ) No difference between groups
Rogind, 1998 <sup>84</sup>	General fitness: LE & low back exercises—ROM, balance, coordination, and muscle strengthening (hip abductors, adductors; quadriceps, hamstrings; gluteal, erector spinae), 3-mo supervised by PT 2 d/wk plus 4 d/wk home program	Control/n = 149 Age: 69 ± 6 Knee OA exercise/n = 11 Age: 69.3 ± 8.2 Knee OA control/n = 12 Age: 73.0 ± 6.5	Physio: Quadriceps isokinetic strength 30° thru 180°/sec Isometric with knee at 90° Postural sway	Both exercise groups reported significantly less pain than controls Significant ↑ (20%) extension strength in exercise groups least affected knee at 30°/sec at 3 mos ( $P < 0.05$ ) ↑ (21%) extension strength in exercise groups least affected knee at 3 mos ( $P = 0.06$ ) No difference between groups

(continued)

Table 2. Continued

Author, Year	Exercises (Modes, Volume, Frequency, Intensity)	Sample Population (Groups/Subjects [n], Age [yr], Disease Classification)	Outcome Measures (Physiologic, Psychological, Functional, Disease Activity)	Results
van Baar, 1998 <sup>76</sup>	Individualized PT exercise program, 12 wk, 1–3/wk, 30-min sessions; intensity: individual	Exercise/n = 93 Age: 68.3 ± 8.4 Control n = 98 Age: 67.7 ± 9.2	Fxn: 20-m walk test  Timed balance test AFI score  Dz: knee effusions  Pain scale  Physio: isometric hip strength  Fxn: disability analyzed on videotapes of performance of specific tasks Dz: Pain (visual analog scale)  No. of medications	Significant ↑ walk velocity in exercise group at 1 yr ( $P < 0.05$ )  No difference between groups Significant ↓ in exercise group at 3 mos and 1 yr ( $P < 0.05$ ); no change in control group Significant ↑ in exercise group at 1 yr (compared with control); significant ↓ in control group at 1 yr (compared with exercisers) Significant ↓ night pain in exercise group at 1 yr ( $P < 0.01$ ) ↑ Hip strength in exercise group ( $P = 0.03$ ) ↓ Disability in exercise group ( $P = 0.04$ ) ↓ Pain in exercise group during last wk and last mo ( $P < 0.01$ for both) ↓ Use of paracetamol in exercise group ( $P = 0.02$ )

NOTE: ↓ = decreased; ↑ = increased; AFI = Algofunctional Index; AIMS = Arthritis Impact Measurement Scales; Dz = disease activity; Fxn = functional measures; HR max = age-predicted heart rate maximum; LE = lower extremities; NR = not reported; OA = osteoarthritis; PT = physical therapy; ROM = range of motion; UE = upper extremities; VO<sub>2</sub>max = maximal aerobic capacity (measurement of aerobic fitness).

SOURCE: Adapted with permission from O'Grady M, Fletcher J, Ortiz S. Therapeutic and physical fitness exercise prescription for older adults with joint disease: an evidence based approach. *Rheum Dis Clin North Am* 2000;26:617–646.

**Table 3. Thermal Modalities: General Physiologic Effects & Precautions**

	Heat	Cold
<b>Physiologic Effects</b>		
Hemodynamic	Vasodilation	Vasoconstriction
Neuromuscular		
Nerve Conduction Velocity	Increased	Decreased
Connective Tissue		
Collagenase Activity	Increased	Decreased
Tendon Extensibility	Increased	Decreased
<b>Clinical Effects</b>		
Pain	Decreased	Decreased
Hemodynamics		
Bleeding	Increased	Decreased
Edema	Increased	Decreased
Inflammation		
Acute	Increased	Decreased
Chronic	Increased	Decreased
Neuromuscular		
Muscle Relaxation	Increased	Increased
Connective Tissue		
Joint Stiffness	Decreased	Increased
<b>Precautions</b>		
Acute Trauma & Inflammation	Contraindicated	Indicated
Impaired Circulation	Contraindicated	Contraindicated
Impaired Sensation	Contraindicated	Contraindicated
Cognitive Impairment	Contraindicated	Contraindicated

tion for developing an appropriate exercise prescription for each patient. Assessment objectives can be divided into two broad categories: arthritis-related factors (current medications, joint pain, inflammation, stability, and range of motion) and impairments associated with inactivity (altered body composition, muscle weakness, and poor cardiovascular fitness). The assessment should include a search for any subclinical or undetected health problems or conditions that could be exacerbated by exercise. In addition, reviewing the patient's expectations along with his or her financial and social resources may improve long-term adherence.<sup>112-116</sup> When all these factors are considered, an exercise prescription can be offered that accommodates to the specific needs and circumstances of the patient.

**The Need For Graded Exercise Testing**

As many older adults may have cardiovascular disease, a complete history and physical examination are needed before prescribing increased physical activity. Contraindications to exercise are presented in Table 4; in general, they are not different from those applicable to younger, healthier adults.<sup>117</sup> Opinions differ regarding the need for a physician-supervised exercise stress test. Cardiovascular response to exercise should be considered for patients with significant risk factors. Such testing assesses cardiac response to exercise and helps to establish an individual's initial aerobic exercise prescription.<sup>118</sup> False positives do occur with exercise stress testing, and there are no consensus recommendations concerning the need to obtain this costly and inconvenient test in older adults who do not have significant cardiovascular disease risk factors. In a review of 14 studies (11 different training protocols) examining the effects of high-intensity strength training in older

adults, only three included exercise stress test as part of their screening procedures.<sup>10,14,16,28,29,119-127</sup> In the largest of these studies, 70 subjects were screened with exercise stress testing; five subjects were excluded because of a resting blood pressure > 160/100, and eight were excluded because of positive exercise stress test results.<sup>28,29</sup>

Although serious cardiovascular events can occur with physical exertion, these usually occur during high-intensity activities. This risk should be considered in light of the fact that regular physical activity of moderate intensity lowers the risk of mortality from cardiovascular disease and can be safely implemented in patients with a low risk for such events.

**How to Start**

The first step in designing an exercise program for the OA patient is to understand which functional problems are most important to the patient. Once disabilities have been

**Table 4. Contraindications to Exercise by the Osteoarthritis Patient**

Absolute	Relative
Uncontrolled arrhythmias	Cardiomyopathy
Third degree heart block	Valvular heart disease
Recent electrocardiographic changes	Poorly controlled blood pressure
Unstable angina	Uncontrolled metabolic disease
Acute myocardial infarction	
Acute congestive heart failure	

inventoried and prioritized, the patient and clinician can set specific short- and long-term goals, which will determine the exercises to be prescribed. Involving the patient in the process enhances long-term adherence.<sup>112-116</sup>

Initially, the program should involve exercises that address the impairments (pain, limited joint range of motion, or muscle weakness) contributing to functional problems. As soon as these impairments begin to improve, a generalized fitness program designed to improve health and functional capacity should begin. Shortly after therapeutic exercise is initiated, fitness training can begin and continue in parallel. The clinician should reinforce the goals and benefits of exercise and familiarize the patient with the specifics of the exercise prescription (intensity, volume, and frequency) and precautions. The latter might include, for example, warnings that physical performance and disease activity can vary from day to day, and that signs of excessive exercise stress include joint pain during activity, pain lasting more than 1 to 2 hours after exercise, swelling, fatigue, and weakness. Patients who are aware of their body's response to exercise and equipped to adjust their training program to avoid immobility may have better long-term adherence with a physical activity program.

#### Basic Exercise Principles and Prescription Components

The basic components for any physical activity program are exercises to improve flexibility, strength, and endurance. Table 5 presents basic recommendations. The training parameters should be individualized for each patient, but all programs are based on general guidelines, as follows.

All exercise prescriptions aimed at improving joint flexibility, muscle strength, or endurance are based on the overload principle: when musculoskeletal tissues are subjected to unaccustomed physiologic stresses, they will adapt and increase their capacity. Overload can be accomplished by increasing the exercise intensity, volume, or frequency, or a combination of these factors.<sup>42</sup>

To be most useful and clear, each exercise prescription specifies exercise intensity, volume, frequency, and progression (Table 5).

- *Intensity* defines the amount of muscular effort or exertion put forth during the activity. The intensity of an activity is typically expressed as a percentage of the individual's maximal capacity. Traditionally, the intensity specified in an exercise program is submaximal (i.e., at levels below the individual's full capacity).
- *Volume* describes how long the exercise is to be performed. For endurance training, volume may be expressed as the amount of time (in number of minutes per exercise session or accumulated minutes per week) the person is engaged in aerobic exercises. For resistance training, volume may be expressed as the number of sets and number of repetitions per set to be performed.
- *Frequency* may be expressed as the number of exercise sessions per week.
- *Progression*, or the gradual application of the overload principle as adaptation occurs, depends on the individual's response to exercise. Although the initial time needed for adaptation to the stress of exercise has not been identified, the range may be 2 to 3 months for most older arthritic adults with reduced physiologic reserve. Progression can be manipulated by changing the intensity, volume, or frequency of training.

The greatest amount of force that a muscle or group of muscles can generate defines strength. A variety of methods have been developed to measure strength. The most commonly used strength measurement is the one repetition maximum or 1RM, defined as the maximum amount of resistance that can be lifted through a full range of motion only once. Typically, the intensity of a strength training program is expressed as a percentage of 1RM. The amount of strength gain depends on the individual's initial level of

**Table 5. Training Parameters: General Guidelines**

Exercise	Intensity	Volume	Frequency
Flexibility: static stretching			
Initial	Stretch to subjective sensation of resistance	1 stretch/key muscle group; hold position 5-15 sec	Once daily
Goal	Stretch to full range of motion	3-5 stretches/key muscle group; hold position 20-30 sec	3-5/wk
Strength: resistance			
Isometric	Low-moderate: 40%-60% MCV	1-10 submaximal contractions involving key muscle group; hold contraction 1-6 sec	Daily
Isotonic	Low: 40% 1 RM Mod: 40%-60% 1 RM High: > 60% 1 RM	10-15 repetitions 8-10 repetitions 6-8 repetitions	2-3/wk
Endurance: aerobic	Low-Mod: 40%-60% of $VO_2\max/HR_{\max}$ RPE: 12-14 = 60%-65% $VO_2\max$ Talk test	Accumulation of 20-30 min/day	3-5/wk

NOTE: 1 RM = one repetition maximum (measurement of isotonic or dynamic strength); MCV = maximal voluntary contraction (measurement of isometric strength); RPE = rating of perceived exertion; HR max = age-predicted heart rate maximum;  $VO_2\max$  = maximal aerobic capacity (measurement of aerobic fitness).

strength and potential and on the training intensity, frequency, and volume.<sup>128,129</sup>

The type of muscle contraction, static or dynamic, differentiates training techniques. A *static* or *isometric* contraction does not change muscle length or move a joint. Isometric strength training occurs when the force of the muscle cannot overcome the applied external resistance (i.e., holding a heavy tray). Strength increases occur primarily at the angle where the muscle was trained, with less improvement at other angles. This drawback limits the usefulness of isometric exercise as the sole form of strength training.

*Dynamic* training is more useful for the person with OA. A *dynamic* contraction both changes muscle length and moves the joint. Dynamic contractions are further classified as isotonic or isokinetic. *Isokinetic* muscle contractions are performed on sophisticated machines that apply variable resistance throughout the range of motion. Isokinetic training, which has been studied in OA patients,<sup>130-132</sup> shows no significant advantages over isotonic strengthening programs. Therefore, from a practical standpoint, isotonic is the recommended form of dynamic strength training for OA patients. An *isotonic* muscle contraction is characterized by variable joint speed exerted against a constant resistance (i.e., free-weight bench press exercise). Isotonic exercise closely corresponds to everyday activities, and strengthening isotonic muscle contractions therefore are recommended for OA patients.

All exercise sessions should have three phases, each of which is essential for reducing the potential for injury and maximizing benefit. The first phase is a warm-up period involving repetitive low-intensity range-of-motion exercises; warm-up lasts 5 to 10 minutes. This phase is important because a proper warm-up prepares the body for more vigorous activity. The second phase is the training period, which provides the overload stimulus to increase joint range of motion, muscle strength, or aerobic capacity, or a combination of these. The final phase, cool-down, lasts 5 minutes and typically involves static stretching of the muscles.

Exercise and other non-pharmacologic interventions are used in parallel with medications to reduce pain and improve function in the older OA patient. The management of symptomatic OA should be adjusted to the needs of the individual patient; an algorithmic approach, though limited, nonetheless helps to organize this complex process into a series of steps (Figure 1). The algorithm highlights the importance of modifiable risk factors in the design of treatment plans that accommodate the heterogeneity of the older OA population and yet facilitates simultaneous implementation of several therapeutic interventions. Such an approach helps to reduce the latency for reducing pain and improving function in older symptomatic OA patients.

## FLEXIBILITY (RANGE-OF-MOTION) EXERCISES

### General Principles

Joint mobility is important to health and to maximal joint range of motion, enhanced muscle performance,<sup>95</sup> reduced risk for injury,<sup>95,133</sup> and improved cartilage nutrition.<sup>134</sup> Flexibility exercises, typically the first step when beginning an exercise program,<sup>34,104</sup> increase the length and elasticity of muscles and periarticular tissues.<sup>135</sup> For the OA patient,

the objectives of such exercises are to decrease stiffness, increase joint mobility, and prevent soft-tissue contractures. Flexibility exercises are often done during the warm-up period or in conjunction with resistance or aerobic activities.

To improve joint range of motion in the OA patient, static stretching is recommended. This stretching technique moves muscles, joints, and periarticular tissues through a range of motion that is comfortable for the patient but that produces some resistances to further movement. Joints, especially those that are painful, should not be over stretched (i.e., stretched to a point that elicits pain), as this may compromise stability. All movement should be through the fullest possible pain-free range. The application of heat prior to stretching may help reduce pain and increase motion.

According to the American College of Sports and Medicine (ACSM), a flexibility program can begin with one stretching exercise per muscle group and should be performed at least 3 times per week.<sup>136</sup> With improvement, the number of repetitions per muscle group can be gradually increased to 4 to 10 repetitions.<sup>137</sup> This general static stretching program should involve the major muscle and tendon groups in the upper and lower extremities (Table 6).

### Static Stretching Exercise: General Recommendations

- Exercise daily when pain and stiffness are minimal (i.e., prior to bedtime).
- Exercises can be preceded by a warm shower or by application of superficial moist heat.
- Relax before beginning stretching exercises.
- Perform movements slowly and extend the range of motion that is both comfortable and produces a slight subjective sensation of resistance. Breathe during each stretch.
- Hold this terminal stretch position for 10 to 30 seconds before slowly returning the joint or muscle group to the resting length.
- Modify the stretching exercises to avoid pain or when the joint is inflamed (decrease the extent of joint range of motion or the duration of holding the static position).

## STRENGTH TRAINING

### General Principles

Strength, an important factor in the performance of daily activities, is an important part of a comprehensive rehabilitation program for the older adult with OA. The aging process, burdens of chronic disease, malnutrition, and inactivity due to OA pain<sup>138</sup> all contribute to reduced muscle mass (sarcopenia) and weakness.<sup>90,139</sup> Studies have shown that resistance training reverses many age-related physiologic changes and can improve function.<sup>94,140-142</sup> The objectives of strength training are to increase the strength of muscles that support the affected joints. The strength training of the individual OA patient should be based on the following principles:

- Specific exercises should be selected on the basis of the patient's joint stability and degree of pain and inflammation.
- Muscles should not be exercised to fatigue.
- Exercise resistance must be submaximal.




Figure 1. Steps in managing osteoarthritis in the older patient.

- Inflamed joints should be isometrically strengthened and involved in only a few repetitions; movements should not be resisted.
- Joint pain lasting 1 hour after exercise and joint swelling indicate excessive activity.

## Isometric Strengthening

### General Principles

Isometric strengthening is indicated when joints are acutely inflamed or unstable. Isometric contractions produce low articular pressures<sup>83</sup> and are well tolerated by OA patients with swollen, painful joints. These exercises can improve muscle strength and static endurance. They prepare the joint for more dynamic movements and are, therefore, typical starting points for most strengthening programs.

Data indicate that strength increases occur when isometric contractions are performed at the muscles' resting length.<sup>143,144</sup> Strength improvements occur primarily at the angle the muscle was trained, with less improvement at different angles<sup>145-147</sup> which hinders the usefulness of this exercise form if the goal is to improve overall function. As

joint instability and pain decrease, the patient's exercise program should gradually shift to dynamic (isotonic) training, as these muscle contractions are used during the performance of activities of daily living.

### Isometric Strength Training Recommendations

- *Exercises:* Include exercises that involve the major muscle groups presented in Table 6.
- *Intensity:* Introductory, isometric contractions should be performed at low intensity. To establish the exercise intensity, ask the patient to maximally contract the muscles targeted for strengthening.<sup>42,129</sup> This is the patient's maximal voluntary contraction and initial training intensity should begin at approximately 30% of this maximal effort. As tolerated by the patient, the intensity should gradually increase to 75% of the maximal voluntary contraction.
- *Volume:* The contraction should be held for no longer than 6 seconds. Initially, one contraction per muscle group should be performed, and the number of repetitions should be gradually increased to eight to 10, as tolerated by the patient.

**Table 6. Key Muscle Groups Targeted for Stretching and Strengthening Exercises**

Head, neck	Extensors, flexors
Shoulder	Forward flexion, extension, abduction, adduction External and internal rotators Scapular retractors and depressors
Elbow	Extensors, flexors
Forearm, wrist	Pronators, supinators Wrist extension, flexors
Hand	Finger flexor, extensors Thumb adductor, abductors
Trunk, low back	Forward flexion, extension, side bending, rotation
Hips	Forward flexion, extension, abduction, adduction External rotation, internal rotation
Knees	Extensors, flexors
Ankle, foot	Dorsiflexors, plantar flexors Inverters, everter Toe flexors, extensors

- The patient should be instructed to breathe during each contraction. Twenty seconds of rest between contractions is suggested.<sup>67,127,143</sup>
- *Frequency:* Exercises should be performed twice daily during acute inflammatory periods. The number of these exercises should be gradually increased to five to 10 times per day, as tolerated by the patient.
- *Progression:* Initially, contractions should be performed at muscle lengths tolerable to the patient. As pain and inflammation decrease, contractions should be performed at different muscle lengths and joint angles.<sup>145</sup> As strength develops, resistance may be added (i.e., contractions against an immovable weight).
- *Precaution:* Contraction > 10 seconds can increase blood pressure.<sup>148</sup>

## Isotonic Training

### General Principles

Isotonic muscle contractions are used to perform activities of daily living. Isotonic strength training has been shown to produce positive effects on energy metabolism,<sup>139</sup> insulin action,<sup>111</sup> bone density,<sup>18</sup> and functional status<sup>18,20,21,141</sup> in healthy older adults. In the absence of inflammation and joint instability, this exercise form is well tolerated by OA patients. Recently, the ACSM published isotonic strength training guidelines. Their recommendations, based on scientific research, are the basis of the exercise recommendations for OA patients outlined below.

### OA Isotonic Exercise Recommendations

Because older OA sufferers with a sedentary lifestyle are likely to have diminished physiologic reserve these exercises should *not* proceed to muscle fatigue.

- *Exercises:* Resistance training should involve eight to 10 exercises involving the major muscle groups.

- *Intensity:* Resistance should begin at 40% of the patient's 1RM. Maximum resistance should be 80% 1RM.
- *Volume:* The beginner should complete one set of four to six repetitions. Exercisers should avoid muscle fatigue.
- *Frequency:* The frequency of training should be a maximum of 2 days per week.
- *Progression:* The progression of resistance training intensity and volume should be gradual to allow time for adaptation. A 5% to 10% increase per week in the amount of resistance used for training seems appropriate.

## Strength Training for Symptomatic Knee OA: An Example

For the medically stable or robust older adult with symptomatic knee OA, some basic exercises aimed at improving quadriceps strength are outlined in Table 7. For those patients with a number of medical problems, the clinician should consider referral to an experienced therapist.

## AEROBIC TRAINING

### General Principles

Aerobic exercise has numerous physiologic benefits that alleviate the deteriorations of aging. These include improved maximal aerobic capacity (measurement of aerobic fitness),<sup>14,15,126,149</sup> insulin action,<sup>150</sup> body composition,<sup>151</sup> and plasma lipoprotein lipid profiles.<sup>75,76,152</sup> Regular aerobic exercise also reduces blood pressure.<sup>153</sup> Meredith and colleagues<sup>138</sup> showed that moderate-intensity training (70% of maximal heart rate), performed 45 minutes per day, 3 days per week for 3 months produced similar aerobic gains in both sedentary young and older adults. The mechanisms for adaptation to aerobic conditioning differ between young and old adults, but improvement in skeletal muscle oxidative capacity and glycogen store are more prominent in the older adult.

The ACSM has set forth standards for the quantity and quality of exercises for developing and improving cardiovascular fitness in an older population.<sup>137</sup> The overload principle is applied to improve an individual's aerobic fitness (VO<sub>2</sub>max). (This principle states that the intensity, frequency, and volume of aerobic exercise must be greater than normal daily activities.) The ACSM recommends that aerobic activities should involve dynamic repetitive movements of large muscle groups.

### Aerobic Exercise Recommendations

- *Exercises:* Activity selection depends on several factors: the patient's current disease activity, joint stability, and resources and interests. The patient should choose a variety of exercise options, to prevent overuse of specific joints and to avoid exercise boredom. Examples of aerobic exercise are bicycling, swimming, low-impact aerobics (i.e., walking, dance, or Tai Chi), or exercising on equipment such as treadmills or rowing machines. Other more utilitarian activities, such as walking the dog, mowing the lawn, raking leaves, or playing golf, are also considered aerobic exercise and should be encouraged. Aquatic exercise is a good choice for OA pa-

Table 7. Home-Based Exercises Designed to Improve Knee Extensor Strength

Exercise	Purpose	Position	Action	Volume and Frequency
Gluteal squeezes	Strengthen buttocks muscles	Lying, sitting, standing	Squeeze buttocks muscles tightly; hold squeeze for 6-7 sec, then slowly relax; rest 2-3 sec between squeezes	Perform 5-7 squeezes 3-5 times a day (before getting out of bed, before climbing stairs, and before going to sleep at night)
Quad set	Strengthen thigh muscles	Lying or standing with legs straight	Squeeze the muscles on top of thigh (this forces the knee cap to move toward the hip); hold squeeze for 6-7 sec, then slowly relax; rest 2-3 sec between squeezes	Perform 5-7 squeezes 3-5 times a day
Short-arc quad set	Strengthen thigh muscle nearest the knee cap	Lying	Place a firm pillow under the knee so that knee is bent; slowly lift foot until leg is straight; hold leg straight for 6-7 sec, then slowly lower leg; rest 2-3 sec between motions	Perform motion 5-7 times 3-5 times a day
Long-arc quad set	Strengthen entire thigh muscle	Sitting	Squeeze muscles on top of thigh to lift foot and straighten leg; hold leg straight for 6-7 sec, then slowly lower leg; rest 2-3 sec between motions	Perform motion 5-7 times 3-5 times a day
Closed-chain, short-arc knee extension	Strengthen hip and thigh muscle	Standing and holding onto a solid object (eg, kitchen counter) for balance	Keeping the back straight, slowly bend knees to 30 degrees, then slowly push back up, extending knees; rest 2-3 sec between motions; as strength improves, increase speed of the motion and depth of knee bend	Perform motion 5-7 times 3-5 times a day

NOTE: Never hold breath during any exercises.

tients; pool exercises performed in warm water (86 °F) provide analgesia for painful muscles and joints. Moreover, the buoyancy of the aquatic environment reduces joint loading, enhances pain-free motion, and provides resistance for strengthening muscle groups around arthritic joints. In addition, pool therapy is commonly a group activity that may help reduce a patient's depression and feelings of isolation. High-impact aerobic training involves rapid application of loads across joint structures and should be avoided, as recent research suggests that the magnitude of joint loading may not be as important in producing pain or damage as the rate of joint loading.<sup>42</sup>

- **Intensity:** Several valid tools are useful for selecting an appropriate exercise intensity, the gold-standard being maximal aerobic power ( $VO_{2MAX}$ ). However, establishing a patient's  $VO_{2MAX}$  is costly and sometimes difficult to obtain. Practical tools that can be helpful in determining appropriate exercise intensity include maximal heart rate ( $HR_{MAX}$ : 220 – (minus sign) age in years), rating of perceived exertion (RPE: a 15-point ordinal scale, 6 to 20), or the “talk test” (whether an exerciser can converse comfortably during the activity without getting short of breath).<sup>154,155</sup> Exercise intensity is considered low to moderate when 1)  $HR_{MAX}$  is between 50% and 75% (i.e. an 80yr old's  $HR_{MAX}$  would be 220–80 or 140 with 50–75% being 70–105), 2) an RPE between 10 and 13, and 3) a positive “talk test.” The aerobic exercise intensity should then range between  $HR_{MAX}$  50% to 60%, RPE 10 and 12, or positive on the “talk test.” For many OA patients, especially those taking medications that control heart rate, the “talk test” or RPE is the simplest method for determining an exercise intensity.
- **Volume:** The recommended volume for the beginner is a minimum of 20 to 30 minutes per day. Some older, sedentary adults are unable to complete 20 to 30 minutes of continuous aerobic activity at low to moderate intensity. An acceptable alternative is four to five shorter exercise bouts (each, a minimum of 5 minutes) performed at slightly higher intensities (i.e., 55% to 60%  $HR_{MAX}$ ) throughout the day.<sup>156,157</sup> Accumulating between 60 and 90 minutes of moderate level physical activity over the course of a week has been included in recent recommendations from the ACSM. As fitness improves, exercise bouts can be lengthened gradually to 20 to 30 minutes of continuous aerobic activity.
- **Frequency:** The initial frequency of training should be at least 3 days but no more than 4 days per week. Frequency of five times per week is not recommended because of increased risk for injury.
- **Progression:** The progression of aerobic training intensity and volume should be gradual to allow time for adaptation (i.e., 2 to 3 months). Following this initial phase of aerobic training, a 2.5% increase per week in the intensity or volume may be compatible with the reduced physiologic reserve associated with older arthritis patients.
- **Precautions:** Musculoskeletal injuries are preventable. More often than not, injuries can be avoided if

the patient gradually works up to the desired activity level and avoids excessive amounts of activity.

### Pharmacologic Therapy

Most authorities agree that the treatment for OA pain should be comprehensive, including both non-pharmacologic and, when necessary, pharmacologic approaches.<sup>1</sup> They also agree that non-pharmacologic therapy should be considered the initial treatment and that pharmacologic agents, such as analgesics and NSAIDs, should be used as adjunctive therapy.<sup>158</sup> Drug therapy for the treatment of OA pain is most effective when used in conjunction with a coordinated program encompassing appropriate non-pharmacologic strategies.<sup>1</sup> Primary physicians who are prescribing exercise for OA patients are encouraged to regularly review the literature on pain management for up-to-date information on the pharmacologic management of pain.

Acetaminophen should be considered the preferred first-line pharmacologic treatment for mild to moderate pain of OA.<sup>1,159–161</sup> Acetaminophen has been shown to provide pain relief comparable to that achieved with NSAIDs,<sup>162,163</sup> without the potential for the gastrointestinal (GI) side effects associated with the use of NSAIDs.<sup>164,165</sup> The daily dosage of acetaminophen should not exceed 4 grams per day.

As an alternative to acetaminophen, a trial of an NSAID (available over the counter [OTC] or by prescription) might be of benefit.<sup>166–171</sup> There is also considerable risk of drug-drug interactions and drug-disease interactions (e.g., congestive heart failure, hypertension, and hepatic and renal disease) with the NSAIDs. Physicians treating OA patients therefore need to take detailed medication histories, including questions about OTC medication use, in order to provide optimal care and recommendations. It has been reported that adverse events with nonselective NSAIDs are more frequent than with any other drug class.<sup>172</sup> It is important to remember that elderly persons are at high risk for side effects of NSAIDs, including GI, platelet, and nephrotoxic effects. Accordingly, NSAIDs should not be used in high doses for long periods of time.<sup>1</sup>

If a patient has a history of gastroduodenal ulcers or of GI bleeding, or develops GI symptoms, one of the new cyclooxygenase (COX-2) inhibitors or selective NSAIDs should be considered.<sup>173,174</sup> The new COX-2 inhibitors have been demonstrated to be as effective as traditional NSAIDs in the management of OA pain. The two currently available COX-2 inhibitors, celecoxib and rofecoxib, have been shown to be as effective as nonselective NSAIDs for mild-to-moderate pain in patients with knee or hip OA.<sup>175,176</sup> However, caution is advised in prescribing a COX-2 (as well as other NSAIDs), as there is the potential for renal complications. There is evidence that rofecoxib tends to cause fluid retention in older adults and that, in addition, taking it without aspirin carries increased cardiovascular risk in this population.<sup>177,178</sup> The choice of agents for treating OA patients with preexisting renal insufficiency requires careful consideration.<sup>179,180</sup>

For patients with OA of the knee and other joints who have mild to moderate pain, topical formulations of analgesics or counterirritants (e.g., methyl salicylate or capsaicin cream, menthol) might be beneficial.<sup>181–184</sup> Expert geriatricians have indicated that pharmacologic modalities for OA pain, besides acetaminophen, NSAIDs, and opioids,

include topical formulations of these agents and intraarticular injections of corticosteroids or hyaluronic acid.<sup>160</sup>

Intraarticular therapy is an alternate approach to pain management in those individuals who either have not obtained relief through systemic medications or in whom oral NSAIDs are contraindicated. This is especially true for patients with OA of the knee. Intraarticular administration of glucocorticoids (e.g., triamcinolone hexacetonide) has been beneficial in treating acute episodes of pain, especially when there is evidence of inflammation and joint effusion.<sup>185,186</sup> More recently, the intraarticular administration of hyaluronic acid preparations has been shown to have efficacy in relieving pain that is not adequately relieved with non-invasive, non-pharmacologic and pharmacologic therapies.<sup>187-189</sup> Other agents that have shown benefit in treating the pain associated with OA of the knee include glucosamine and chondroitin.<sup>190,191</sup> However, additional studies are necessary to demonstrate long-term safety and efficacy of these agents.

For some patients with severe OA pain that is refractory to other forms of therapy, stronger analgesic drugs may be required. Carefully titrated opioid analgesic drugs may be preferable to NSAIDs, cortisone, or other pharmacologic or invasive interventions that pose appreciable risks in older people.<sup>1</sup> It has also been suggested that opioid analgesics may be better for treating acute exacerbations of OA pain than for long-term use.<sup>159</sup>

## CONCLUSIONS

Chronic pain and functional dependency are not inevitable consequences of aging with joint disease. Identifying the modifiable risk factors (Table 1) enables practitioners to focus their therapeutic interventions, reduce pain, and improve function. Randomized, controlled trials clearly show that regular moderate-level exercise does not exacerbate OA pain or accelerate the pathological process of OA. Furthermore, these studies strongly indicate that increasing the level of physical activity in OA patients reduces pain and morbidity. Exercise programs should be individualized to address the specific needs of the patient. The goals of any exercise program should focus on controlling pain, increasing flexibility, and improving muscle strength and endurance.

## ACKNOWLEDGMENTS

The AGS Panel on Osteoarthritis and Exercise was chaired by Paul Katz, MD, University of Rochester, Rochester, New York.

On behalf of the Panel, Michael O'Grady, MD, Emory University School of Medicine, Atlanta, Georgia, was the primary author of these consensus recommendations.

The members of the panel were Gail Davis, EdD, RN, Texas Women's University, Denton, Texas; Carlos H. Rojas-Fernandez, PharmD, Texas Tech University Health Sciences Center, School of Pharmacy, Amarillo, Texas; Bruce Ferrell, MD, UCLA School of Medicine, Los Angeles, California. Roger Levy, MD, Hospital for Special Surgery, New York, New York; David C. Neiman, DrPH, Appalachian State University, Boone, North Carolina; Mark A. Young, MD, FACP, Maryland Rehabilitation Center, Johns Hopkins University School of Medicine, Baltimore, Maryland.

Research services were provided by Sue Radcliff, Independent Researcher, Denver, CO, USA. Editorial services were provided by Barbara B. Reitt, PhD, ELS(D) of Reitt Editing Services. Additional research and administrative support was provided by Adrienne Prassas and Nancy Lundberg, Professional Education and Publications, American Geriatrics Society, New York, NY, USA.

The following organizations with special interest and expertise in the management of osteoarthritis and exercise in older person provided expert review of an earlier version of the guidelines: American Academy of Family Physicians, American Academy of Orthopaedic Surgeons, American Academy of Physical Medicine and Rehabilitation, American Nurses Association, and the American Physical Therapy Association.

Funding was provided as an unrestricted educational grant from McNeil Consumer Health Care, Fort Washington, Pennsylvania.

## REFERENCES

1. AGS Panel on Chronic Pain in Older Persons. The management of chronic pain in older persons: clinical practice guideline. *J Am Geriatr Soc* 1998;46:635-651.
2. United States Census Bureau. Available at URL: <http://www.census.gov> 1999
3. Health-related quality of life among adults with arthritis - behavioral risk factor surveillance system, 11 states, 1996-1998. *MMWR Morb Mortal Wkly Rep* 2000;49:366-369.
4. Olshansky SJ, Carnes BA, Cassel CK. The aging of the human species. *Sci Am* 1993;268:46-52.
5. Verbrugge LM, Lepkowski JM, Konkol LL. Levels of disability among U.S. adults with arthritis. *J Gerontol* 1991;46:S71-S83.
6. Mazzuca SA, Brandt KD, Katz BP et al. Therapeutic strategies distinguish community based primary care physicians from rheumatologists in the management of osteoarthritis. *J Rheumatol* 1993;20:80-86.
7. Prevalence and impact of arthritis among women - United States, 1989-1991. *MMWR Morb Mortal Wkly Rep* 1995;44:331-335.
8. Michet CJ. Osteoarthritis. *Prim Care* 1993;20:815-826.
9. Lawrence RC, Helmick CG, Arnett FC et al. Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arth Rheum* 1998;41:778-799.
10. Frontera WR, Meredith CN, O'Reilly KP et al. Strength conditioning in older men: skeletal muscle hypertrophy and improved function. *J Appl Physiol* 1988;64:1038-1044.
11. Loeser RF Jr. Aging and the etiopathogenesis and treatment of osteoarthritis. *Rheum Dis Clin North Am* 2000;26:547-567.
12. McAuley E, Katula J. Physical activity interventions in the elderly: influence on physical health and psychological function. *Annu Rev of Gerontol Geriatr* 1998;18:111-154.
13. Evans WJ. What is sarcopenia? *J Gerontol A Biol Sci Med Sci* 1995;50:5-8.
14. Fiatarone MA, Marks EC, Ryan ND et al. High-intensity strength training in nonagenarians. Effects on skeletal muscle. *JAMA* 1990;263:3029-3034.
15. Fiatarone MA, Evans WJ. The etiology and reversibility of muscle dysfunction in the aged. *J Gerontol* 1993;48:77-83.
16. Fiatarone MA, O'Neill EF, Ryan ND et al. Exercise training and nutritional supplements for physical frailty in very elderly people. *N Engl J Med* 1994;330:1769-1775.
17. Fisher NM, Gresham GE, Abrams M et al. Quantitative effects of physical therapy on muscular and functional performance in subjects with osteoarthritis of the knees. *Arch Phys Med Rehabil* 1993;74:840-847.
18. Fisher NM, Kame VD, Rouse L et al. Quantitative evaluation of a home exercise program on muscle and functional capacity of patients with osteoarthritis. *Am J Phys Med Rehabil* 1994;73:413-420.
19. Fleg JL, Lakatta EG. Role of muscle loss in the age-associated reduction in VO<sub>2</sub>max. *J Appl Physiol* 1988;65:1147-1151.
20. Gardner GW. Specificity of strength changes of the exercised and non-exercised limb following isometric training. *Res Q* 1962;34:98-101.
21. Gardner GW. Effect of isometric and isotonic exercise on joint motion. *Arch Phys Med Rehabil* 1966;47:24-30.
22. Goldberg AP. Aerobic and resistive exercise modify risk factors for coronary heart disease. *Med Sci Sports Exerc* 1989;21:669-674.
23. Graves JE, Pollock ML, Jones AE et al. Specificity of limited range of motion variable resistance training. *Med Sci Sports Exerc* 1989;21:84-89.

24. Grimby G. Muscle performance and structure in the elderly as studied cross-sectionally and longitudinally. *J Gerontol A Biol Sci Med Sci* 1995;50:17-22.
25. Guralnik JM, Branch LG, Cummings SR et al. Physical performance measures in aging research. *J Gerontol* 1989;44:M141-M146.
26. Gussoni M, Margonato V, Ventura R et al. A. Energy costs of walking with hip joint impairments. *Phys Ther* 1990;70:295-301.
27. Guyatt GH, Sullivan MJ, Thompson PJ et al. The 6-minute walk: a new measure of exercise capacity in patients with chronic heart failure. *Can Med Assoc J* 1985;132:919-923.
28. Hagberg JM, Graves JE, Limacher M et al. Cardiovascular responses of 70- to 79-yr-old men and women to exercise training. *J Appl Physiol* 1989;66:2589-2594.
29. Hagberg JM, Montain SJ, Martin WH III et al. Effect of exercise training in 60- to 69-year old persons with essential hypertension. *Am J Cardiol* 1989;64:348-353.
30. Hakkinen A, Hakkinen K, Hannonen P. Effects of strength training on neuromuscular function and disease activity in patients with recent-onset inflammatory arthritis. *Scand J Rheumatol* 1994;23:237-242.
31. Hakkinen A, Malkia E, Hakkinen K et al. Effects of detraining subsequent to strength training on neuromuscular function in patients with inflammatory arthritis. *Br J Rheumatol* 1997;36:1075-1081.
32. Hansen TM, Hansen G, Langgaard AM et al. Long term physical training in rheumatoid arthritis. A randomized trial with different training programs and blinded observers. *Scand J Rheumatol* 1993;22:107-112.
33. Harkcom TM, Lampman RM, Banwell BF et al. Therapeutic value of graded aerobic exercise training in rheumatoid arthritis. *Arthritis Rheum* 1985;28:32-39.
34. Paffenbarger RS, Hyde RT, Wing AL et al. Physical activity and all-cause mortality, and longevity of college alumni. *N Engl J Med* 1986;314:605-613.
35. Harries UJ, Bassey EJ. Torque-velocity relationships for the knee extensors in women in their 3rd and 7th decades. *Eur J Appl Physiol Occup Physiol* 1990;60:187-190.
36. Hasselkus BR, Shambes GM. Aging and postural sway in women. *J Gerontol* 1975;30:661-667.
37. Heath G, Hagberg J, Ehsani A et al. A physiological comparison of young and older endurance athletes. *J Appl Physiol* 1981;51:634-640.
38. Hellebrandt FA, Braun GL. The influence of sex and age on the postural sway of man. *Am J Phys Anthropol* 1939;24:347-360.
39. Brandt KD. The importance of nonpharmacologic approaches in management of osteoarthritis. *Am J Med* 1998;105:395-445.
40. Hochberg MC, Altman RD, Brandt KD et al. Guidelines for the medical management of osteoarthritis. Part II. Osteoarthritis of the knee. *Arth Rheum* 1995;38:1541-1546.
41. Bradley LA, Alberts KR. Psychological and behavioral approaches to pain management for patients with rheumatic disease. *Rheum Dis Clin North Am* 1999;25:215-232.
42. O'Grady M, Fletcher J, Ortiz S. Therapeutic and physical fitness exercise prescription for older adults with joint disease: an evidence-based approach. *Rheum Dis Clin North Am* 2000;26:617-646.
43. Schouten JS, Valkenburg HA. Classification criteria: methodological considerations and results from a 12-year following study in the general population. *J Rheumatol* 1995;43:44-45.
44. Oliveria SA, Felson DT, Reed JI et al. Incidence of symptomatic hand, hip, and knee osteoarthritis among patients in a health maintenance organization. *Arth Rheum* 1995;38:1134-1141.
45. Bagge E, Bjelle A, Eden S et al. Osteoarthritis in the elderly: clinical and radiological findings in 79 and 85 year olds. *Ann Rheum Dis* 1991;50:535-539.
46. Lawrence RC, Hochberg MC, Kelsey JL et al. Estimates of the prevalence of selected arthritic and musculoskeletal diseases in the United States. *J Rheumatol* 1989;16:427-441.
47. Creamer P, Hochberg MC. Osteoarthritis. *Lancet* 1997;350:503-509.
48. Panush RS. Physical activity, fitness, and osteoarthritis. In: Bouchard C, Shephard RJ, Stephens T, eds. *Physical Activity, Fitness and Health: International Proceedings and Consensus Statement*. Champaign, IL: Human Kinetics, 1994:712-723.
49. Bagge E, Bjelle A, Eden S et al. A longitudinal study of the occurrence of joint complaints in elderly people. *Age Ageing* 1992;21:160-167.
50. Egger P, Cooper C, Hart DJ et al. Patterns of joint involvement in osteoarthritis of the hand: The Chingford Study. *J Rheumatol* 1995;22:1509-1513.
51. Maurer K. Basic data on arthritis knee, hip and sacroiliac joints in adults ages 25-74 years, United States 1971-1975. National Center for Health Statistics, Department of Health, Education and Welfare. 1979;Series 11:1-31.
52. Lethbridge-Cejku M, Scott WW, Reichle R et al. Association of radiographic features of osteoarthritis of the knee with knee pain: data from the Baltimore Longitudinal Study of Aging. *Arth Care Res* 1995;8:182-188.
53. Felson DT, Anderson JJ, Naimark A et al. Obesity and knee osteoarthritis, The Framingham Study. *Ann Intern Med* 1988;109:18-24.
54. Felson DT, Zhang Y, Hannan MT et al. The incidence and natural history of knee osteoarthritis in the elderly: The Framingham Osteoarthritis Study. *Arth Rheum* 1995;38:1500-1505.
55. van Saase JL, van Romunde LK, Cats A et al. Epidemiology of osteoarthritis: Zoetermeer survey. Comparison of radiological osteoarthritis in a Dutch population with that in 10 other populations. *Ann Rheum Dis* 1989;48:271-280.
56. Felson DT, Naimark A, Anderson JJ et al. The prevalence of knee osteoarthritis in the elderly: The Framingham Osteoarthritis Study. *Arth Rheum* 1987;30:914-918.
57. Verbrugge LM. Women, men and osteoarthritis. *Arth Care Res* 1995;6:212-220.
58. Panush R. Arthritis. In: Neiman DC, ed. *The Exercise-Health Connection*. Champaign, IL: Human Kinetics, 1998:119-131.
59. Felson DT, Zhang Y, Hannan MT et al. Risk factors for incident radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arth Rheum* 1997;40:728-733.
60. Hochberg MC, Lethbridge-Cejku M, Scott WW et al. The association of body weight, body fatness and body fat distribution with osteoarthritis of the knee: data from the Baltimore Longitudinal Study of Aging. *J Rheumatol* 1995;22:488-493.
61. Vingard E, Alfredsson L, Malchau H. Lifestyle factors and hip arthrosis. A case referent study of body mass index, smoking and hormone therapy in 503 Swedish women. *Acta Orthop Scand* 1997;68:216-220.
62. Felson DT, Zhang Y, Anthony JM et al. Weight loss reduces the risk for symptomatic knee osteoarthritis in women. *Ann Intern Med* 1992;116:535-539.
63. Cooper C, Inskip H, Croft P et al. Individual risk factors for hip osteoarthritis: obesity, hip injury, and physical activity. *Am J Epidemiol* 1998;147:516-522.
64. Engelhart M, Kondrup J, Hoie LH et al. Weight reduction in obese patients with rheumatoid arthritis, with preservation of body cell mass and improvement of physical fitness. *Clin Exp Rheumatol* 1996;14:289-293.
65. Hochberg MC, Lethbridge-Cejku M. Epidemiologic considerations in the primary prevention of osteoarthritis. In: Hamerman D, ed. *Osteoarthritis: Public Health Implications for an Aging Population*. Baltimore: Johns Hopkins University Press, 1997:169-186.
66. Toda Y, Toda T, Takemura S et al. Changes in body fat, but not body weight or metabolic correlates of obesity, is related to symptomatic relief of obese patients with knee osteoarthritis after a weight control program. *J Rheumatol* 1998;25:2181-2186.
67. Slemenda CW, Brandt KD, Heilman DK et al. Quadriceps weakness and osteoarthritis of the knees. *Ann Intern Med* 1997;127:97-104.
68. Slemenda CW, Heilman DK, Brandt KD et al. Reduced quadriceps strength relative to body weight: a risk factor for knee osteoarthritis in women? *Arth Rheum* 1998;41:1951-1959.
69. Turner AP, Barlow JH, Heathcote-Elliott C. Long-term health impact of playing professional football in the United Kingdom. *Br J Sports Med* 2000;34:332-336.
70. Felson DT, Zhang Y. An update on the epidemiology of knee and hip osteoarthritis with a view to prevention. *Arth Rheum* 1998;41:1343-1355.
71. Roos H, Lauren M, Adalberth T et al. Knee osteoarthritis after meniscectomy: prevalence of radiographic changes after twenty-one years, compared with matched controls. *Arth Rheum* 1998;41:687-693.
72. Roos H, Adalberth T, Dahlberg L et al. Osteoarthritis of the knee after injury to the anterior cruciate ligament or meniscus: the influence of time and age. *Osteoarth Cartilage* 1995;3:261-267.
73. Gelber AC, Hochberg MC, Mead LA et al. Joint injury in young adults and risk for subsequent knee and hip osteoarthritis. *Ann Intern Med* 2000;133:321-328.
74. Hurley MV. The role of muscle weakness in the pathogenesis of osteoarthritis. *Rheum Dis Clin North Am* 1999;25:283-298.
75. van Baar ME, Dekker J, Oostendorp RA et al. The effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a randomized clinical trial. *J Rheumatol* 1998;25:2432-2439.
76. Ettinger WH, Jr., Burns R, Messier SP et al. A randomized trial comparing aerobic exercise with resistance exercise with a health education program in older adults with knee osteoarthritis: The Fitness Arthritis and Seniors Trial (FAST). *JAMA* 1997;277:25-31.
77. Pai YC, Rymer WZ, Chang RW et al. Effect of age and osteoarthritis on knee proprioception. *Arthritis Rheum* 1997;40:2260-2265.
78. Hurley MV, Scott DL, Rees J et al. Sensorimotor changes and functional performance in patients with knee osteoarthritis. *Ann Rheum Dis* 1997;56:641-648.
79. Sharma L, Pai YC, Holtkamp K et al. Is knee joint proprioception worse in the arthritic knee versus the unaffected knee in unilateral knee osteoarthritis? *Arth Rheum* 1997;40:1518-1525.
80. Hurley MV, Scott DL. Improvements in quadriceps sensorimotor function

- and disability of patients with knee osteoarthritis following a clinically practicable exercise regime. *Br J Rheumatol* 1998;37:1181-1187.
81. Minor MA, Hewett JE, Weibel RR et al. Efficacy of physical conditioning exercise in patients with rheumatoid arthritis and osteoarthritis. *Arth Rheum* 1989;32:1396-1405.
  82. Kovar PA, Allegrante JP, MacKenzie CR et al. Supervised fitness walking in patients with osteoarthritis of the knee. A randomized controlled trial. *Ann Intern Med* 1992;116:529-534.
  83. Rogind H, Bibow-Nielsen B, Jensen B et al. The effects of a physical training program on patients with osteoarthritis of the knees. *Arch Phys Med Rehabil* 1998;79:1421-1427.
  84. Lane NE. Physical activity at leisure and risk of osteoarthritis. *Ann Rheum Dis* 1996;55:682-684.
  85. Fries JF, Singh G, Morfield D et al. Running and the development of disability with age. *Ann Intern Med* 1994;121:502-509.
  86. Buckwalter JA, Lane NE. Athletics and osteoarthritis. *Am J Sports Med* 1997;25:873-881.
  87. Saxon L, Finch C, Bass S. Sports participation, sports injuries and osteoarthritis: implications for prevention. *Sports Med* 1999;28:123-135.
  88. Marti B, Knobloch M, Tschopp A et al. Is excessive running predictive of degenerative hip disease? Controlled study of former elite athletes. *BMJ* 1989;299:91-93.
  89. Croft P, Coggon D, Cruddas M et al. Osteoarthritis of the hip: an occupational disease in farmers. *BMJ* 1992;304:1269-1272.
  90. Puett DW, Griffin MR. Published trials of nonmedicinal and noninvasive therapies for hip and knee osteoarthritis. *Ann Intern Med* 1994;121:133-140.
  91. Minor MA. Exercise in the management of osteoarthritis of the knee and hip. *Arth Care Res* 1994;7:198-204.
  92. Lyngberg KK, Harreby M, Bentzen H et al. Elderly rheumatoid arthritis patients on steroid treatment tolerate physical training without an increase in disease activity. *Arch Phys Med Rehabil* 1994;75:1189-1195.
  93. Coleman EA, Buchner DM, Cress ME et al. The relationship of joint symptoms with exercise performance in older adults. *J Am Geriatr Soc* 1996;44:14-21.
  94. Rall LC, Meydani SN, Kehayias JJ et al. The effect of progressive resistance training in rheumatoid arthritis. Increased strength without changes in energy balance or body composition. *Arth Rheum* 1996;39:415-426.
  95. Noreau L, Moffett H, Drolet M et al. Dance-based exercise program in rheumatoid arthritis. *Am J Phys Med Rehabil* 1997;76:109-113.
  96. Christmas C, Andersen RA. Exercise and older patients: guidelines for the clinician. *J Am Geriatr Soc* 2000;48:318-324.
  97. Messier SP, Thompson CD, Ettinger Jr. WH. Effects of long-term aerobic or weight training regimens on gait in an older osteoarthritic population. *J Appl Biomech* 1997;13:205-225.
  98. Messier SP, Royer TD, Craven TE et al. Long-term exercise and its effect on balance in older, osteoarthritic adults: results from Fitness, Arthritis, and Seniors Trial (FAST). *J Am Geriatr Soc* 2000;48:131-138.
  99. Minor MA. Exercise in the treatment of osteoarthritis. *Rheum Dis Clin North Am* 1999;25:397-415.
  100. Schilke JM, Johnson GO, Housh TJ et al. Effects of muscle-strength training on the functional status of patients with osteoarthritis of the knee joint. *Nurs Res* 1996;45:68-72.
  101. Brandt KD, Heilman DK, Slemenda C et al. Quadriceps strength in women with radiographically progressive osteoarthritis of the knee and those with stable radiographic changes. *J Rheumatol* 1999;26:2431-2437.
  102. Ettinger WH, Jr., Afbale RF. Physical disability from knee osteoarthritis: the role of exercise as an intervention. *Med Sci Sports Exerc* 1994;26:1435-1440.
  103. Minor MA, Brown JD. Exercise maintenance of persons with arthritis after participation in a class experience. *Health Educ Q* 1993;20:83-95.
  104. O'Reilly SC, Muir KR, Doherty M. Effectiveness of home exercise on pain and disability from osteoarthritis of the knee: a randomized controlled trial. *Ann Rheum Dis* 1999;58:15-19.
  105. Hochberg MC, Altman RD, Brandt KD et al. Guidelines for the medical management of osteoarthritis. Part I. Osteoarthritis of the hip. *Arth Rheum* 1995;38:1535-1540.
  106. Weinberger M, Tierney WN, Booher P et al. The impact of increased contact on psychosocial outcomes in patients with osteoarthritis: a randomized controlled trial. *J Rheumatol* 1991;18:849-854.
  107. Hirano PC, Laurent DD, Lorig KR. Arthritis patient education studies, 1987-1991: a review of the literature. *Patient Educ Couns* 1994;24:9-54.
  108. Lorig KR, Mazonson PD, Holman HR. Evidence suggesting that health education for self-management in patients with chronic arthritis has sustained health benefits while reducing health care costs. *Arth Rheum* 1993;36:439-445.
  109. Lorig KR, Holman HR. Arthritis self-management studies: a 12 year review. *Health Educ Q* 1993;20:17-28.
  110. Keefe FJ, Caldwell DS, Baucom D et al. Spouse-assisted coping skills training in the management of osteoarthritic knee pain. *Arth Care Res* 1996;9:279-291.
  111. Stenstrom CH, Lindell B, Swanberg E et al. Intensive dynamic training in water for rheumatoid arthritis functional class II - a long-term study of effects. *Scand J Rheumatol* 1991;20:358-365.
  112. King AC, Rejeski WJ, Buchner DM. Physical activity interventions targeting older adults. A critical review and recommendations. *Am J Prev Med* 1998;15:316-333.
  113. Stewart AL, Mills KM, Sepsis PG et al. Evaluation of CHAMPS, a physical activity promotion program for older adults. *Ann Behav Med* 1998;19:353-361.
  114. King AC, Sallis JF, Dunn AL et al. Overview of the Activity Counseling Trial (ACT) intervention for promoting physical activity in primary health care settings. Activity Counseling Trial Research Group. *Med Sci Sports Exerc* 1998;30:1086-1096.
  115. King AC, Pruitt LA, Phillips W et al. Comparative effects of two physical activity programs on measured and perceived physical functioning and other health-related quality of life outcomes in older adults. *J Gerontol A Biol Sci Med Sci* 2000;55A:M74-M83.
  116. Brownson RC, Eyster AA, King AC et al. Reliability of information on physical activity and other chronic disease risk factors among U.S. women aged 40 years or older. *Am J Epidemiol* 1999;149:379-391.
  117. Nordemar R, Ekblom B, Zachrisson L et al. Physical training in rheumatoid arthritis: a controlled long-term study. *Scand J Rheumatol* 1981;10:17-23.
  118. Gill TM, DiPietro L, Krumholz HM. Role of exercise stress testing and safety monitoring for older persons starting an exercise program. *JAMA* 2000;284:342-349.
  119. Charette SL, McEvoy L, Pyka G et al. Muscle hypertrophy in response to resistance training in older women. *J Appl Physiol* 1991;70:1912-1916.
  120. Koffler KH, Menkes A, Redmond RA et al. Strength training accelerates gastrointestinal transit in middle-aged and older men. *Med Sci Sports Exerc* 1992;24:415-419.
  121. Pratley R, Nicklas B, Rubin M et al. Strength training increases resting metabolic rate and norepinephrine levels in healthy 50- to 65-year-old men. *J Appl Physiol* 1994;76:133-137.
  122. Pyka G, Lindenberg E, Charette S et al. Muscle strength and fiber adaptations to a year-long resistance training program in elderly men and women. *J Gerontol* 1994;49:M22-M27.
  123. Campbell WW, Crim MC, Young VR et al. Increased energy requirements and changes in body composition with resistance training in older adults. *Am J Clin Nutr* 1994;60:167-175.
  124. Nelson ME, Fiatarone MA, Morganti CM et al. Effects of high intensity strength training on multiple risk factors for osteoporotic fractures. A randomized controlled trial. *JAMA* 1994;272:1909-1914.
  125. Morganti CM, Nelson ME, Fiatarone MA et al. Strength improvements with 1 yr of progressive resistance training in older women. *Med Sci Sports Exerc* 1995;27:906-912.
  126. Taaffe DR, Pruitt L, Reim J et al. Effect of sustained resistance training on basal metabolic rate in older women. *J Am Geriatr Soc* 1995;43:465-471.
  127. Singh NA, Clements KM, Fiatarone MA. A randomized controlled trial of progressive resistance training in depressed elders. *J Gerontol A Biol Sci Med Sci* 1997;52:M27-M35.
  128. Liberson WT, Asa MM. Further studies of brief isometric exercise. *Arch Phys Med Rehabil* 1959;40:330-336.
  129. Kraemer WJ, Newton RU. Training for muscle power. *Phys Med Rehabil Clin North Am* 2000;11:341-368.
  130. Kohrt WM, Spina RJ, Holloszy JO et al. Prescribing exercise intensity for older women. *J Am Geriatr Soc* 1998;46:129-133.
  131. Stevenson ET, Davy KP, Seals DR. Hemostatic, metabolic, and androgenic risk factors for coronary heart disease in physically active and less active postmenopausal women. *Arterioscler Thromb Vasc Biol* 1995;15:669-677.
  132. Stratton JR, Levy WC, Cerqueira MD et al. Cardiovascular responses to exercise. Effects of aging and exercise training in healthy men. *Circulation* 1994;89:1648-1655.
  133. Oddis C. New perspectives on osteoarthritis. *Am J Med* 1996;100:105-155.
  134. Ogawa T, Spina RJ, Martin WH, III et al. Effects of aging, sex and physical training on cardiovascular responses to exercise. *Circulation* 1992;86:494-503.
  135. Palmoski MJ, Bolyer RA, Brandt KD. Joint motion in the absence of normal loading does not maintain normal articular cartilage. *Arth Rheum* 1980;23:325-334.
  136. Pollock ML, Mengelkoch LJ, Graves JE et al. Twenty-year follow-up of aerobic power and body composition of older track athletes. *J Appl Physiol* 1997;82:1508-1516.
  137. Lindh M. Increase of muscle strength from isometric quadriceps exercise at different knee angles. *Scand J Rehab Med* 1979;11:33-36.

138. Meredith CN, Frontera WR, Fisher EC et al. Peripheral effects of endurance training in young and old subjects. *J Appl Physiol* 1989;66:2844-2849.
139. Pyykko I, Aalto H, Hytonen M et al. Effect of age on postural control. In: Amlund B, Berthoz A, Clarac F, eds. *Posture and Gait: Development, Adaptation and Modulation*. New York: Elsevier Science, 1988:95-104.
140. Rejeski WJ, Ettinger WH, Jr., Shumaker S et al. Assessing performance-related disability in patients with knee osteoarthritis. *Osteoarth Cartilage* 1995;3:157-167.
141. Robertson RJ, Noble BN. Perception of physical exertion: methods, mediators, and applications. In: Holloszy JO, ed. *Exercise and Sport Sciences Reviews*. Baltimore: Williams & Wilkins, 1997:407-452.
142. Rodeheffer RJ, Gerstenblith G, Becker LC et al. Exercise cardiac output is maintained with advancing age in healthy human subjects: cardiac dilatation and increased stroke volume compensate for a diminished heart rate. *Circulation* 1984;69:203-213.
143. Schwartz RS, Shuman WP, Larson V et al. The effect of intensive endurance exercise training on body fat distribution in young and older men. *Metabolism* 1991;40:545-551.
144. Seals DR, Allen WK, Hurley BF et al. Elevated high-density lipoprotein cholesterol levels in older endurance athletes. *Am J Cardiol* 1984;54:390-393.
145. Seals DR, Hagberg JM, Allen WK et al. Glucose tolerance in young and older athletes and sedentary men. *J Appl Physiol* 1984;56:1521-1525.
146. Seals DR, Hagberg JM, Hurley BF et al. Effects of endurance training on glucose tolerance and plasma lipid levels in older men and women. *JAMA* 1984;252:645-649.
147. Seals DR, Hagberg JM, Hurley BF et al. Endurance training in older men and women. I. Cardiovascular responses to exercise. *J Appl Physiol* 1984;57:1024-1029.
148. Spina RJ, Miller TR, Bogenhagen WH et al. Gender-related differences in left ventricular filling dynamics in older subjects after endurance exercise training. *J Gerontol A Biol Sci Med Sci* 1996;51:B232-B237.
149. Danneskiold-Samsoe BV, Kofod V, Munter J et al. Muscle strength and functional capacity in 78-81 year old men and women. *Eur J Appl Physiol* 1984;52:310-314.
150. Doucette SA, Goble EM. The effect of exercise on patellar tracking in lateral patellar compression syndrome. *Am J Sports Med* 1992;20:434-440.
151. Tinetti ME. Performance-oriented assessment of mobility problems in elderly people. *J Am Geriatr Soc* 1986;34:119-126.
152. Tinetti ME, Baker DI, McAvay G et al. A multifactorial intervention to reduce the risk of falling among elderly people living in the community. *N Engl J Med* 1994;331:821-827.
153. Van Deusen J, Jarlowe D. One-year follow up results of ROM dance research. *Occ Ther J Res* 1988;8:52-54.
154. Ware JE, Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF36): I. Conceptual framework and item selection. *Med Care* 1992;30:473-483.
155. Wessel J, Quinney HA. Pain experienced by persons with rheumatoid arthritis during isometric and isokinetic exercise. *Physiother Can* 1984;36:131-134.
156. Wolf SL, Barnhart HX, Kutner NG et al. Reducing frailty and falls in older persons: an investigation of Tai Chi and computerized balance training. Atlanta FICSIT Group. Frailty and Injuries: Cooperative Studies of Intervention Techniques. *J Am Geriatr Soc* 1996;44:489-497.
157. Worrell TW, Smith TL, Winegardner J. Effect of stretching on hamstring muscle performance. *J Orthop Sports Phys Ther* 1994;20:154-159.
158. Kruger JM, Helmick CG, Callahan LF, Haddix AC. Cost-effectiveness of the arthritis self-help course. *Arch Intern Med* 1998;158:1245-1249.
159. Osteoarthritis: non-narcotic analgesics. *Geriatric Pharmaceutical Care Guidelines*. Omnicare Inc., Covington KY 2000:371-378.
160. Reuben D, Herr K, Pacala J et al. Table 53. Acetaminophen and nonsteroidal anti-inflammatory drugs. In: *Geriatrics at Your Fingertips*. New York: American Geriatrics Society, 2000:88-91.
161. Williams M. Conditions that affect older people: joints, muscles, and bones. In: *The American Geriatrics Society's Complete Guide to Aging and Health*. New York: Harmony Books, 1995:259-260.
162. Bradley JD, Brandt KD, Katz BP et al. Comparison of an anti-inflammatory dose of ibuprofen, an analgesic dose of ibuprofen, and acetaminophen in the treatment of patients with osteoarthritis of the knee. *N Engl J Med* 1991;325:87-91.
163. Eccles M, Freemantle N, Mason J. North of England evidence based guideline development project: summary guideline for non-steroidal anti-inflammatory drugs versus basic analgesia in treating the pain of degenerative arthritis. *BMJ* 1998;317:526-530.
164. Henry D, Lim LL, Garcia Rodriguez LA et al. Variability in risk of gastrointestinal complications with individual non-steroidal anti-inflammatory drugs: results of a collaborative meta-analysis. *BMJ* 1996;312:1563-1566.
165. Phillips AC, Polisson RP, Simon LS. NSAIDs and the elderly: Toxicity and economic implications. *Drugs Aging* 1997;10:119-130.
166. Malmstrom K, Daniels S, Kotey P et al. Comparison of rofecoxib and celecoxib, two cyclooxygenase-2 inhibitors, in postoperative dental pain: a randomized, placebo- and active comparator-controlled clinical trial. *Clin Ther* 1999;21:1653-1663.
167. Simon LS, Weaver AL, Graham DY et al. Anti-inflammatory and upper gastrointestinal effects of celecoxib in rheumatoid arthritis: a randomized controlled trial. *JAMA* 1999;282:1921-1928.
168. Stubanus M, Reigger GAJ, Kammert MC. Renal side-effects of cyclo-oxygenase-2 inhibitor use. Letter to editor. *Lancet* 2000;355:753.
169. Brater DC. Effects of nonsteroidal anti-inflammatory drugs on renal function: focus on cyclooxygenase-2-selective inhibition. *Am J Med* 1999;107:65S-71S.
170. Swan SK, Lasserter KC, Ryan CF et al. Renal effects of multiple-dose rofecoxib (R), a Cox-2 inhibitor, in elderly subjects, abstract A3248. *J Am Soc Nephrol* 1999;10:641A.
171. Helme RD, Gibson SJ. Pain in the elderly. In: Jensen TS, Turner JA, Wiesenfeld-Hallin Z, eds. *Proceedings of the 8th World Congress on Pain: Progress in Pain Research and Management*. Seattle: IASP Press, 1997:919-944.
172. Langman MJ, Jensen DM, Watson DJ et al. Adverse upper gastrointestinal effects of rofecoxib compared with NSAIDs. *JAMA* 1999;282:1929-1933.
173. Simon L. Nonsteroidal anti-inflammatory drugs and their effects. The importance of COX "selectivity". *J Clin Rheumatol* 1996;2:135-140.
174. Crofford LJ, Lipsky PE, Brooks P et al. Basic biology and clinical application of specific cyclooxygenase-2 inhibitors. *Arth Rheum* 2000;43:4-13.
175. Ehrlich EW, Schnitzer TJ, McIlwain H et al. Effect of specific COX-2 inhibition in osteoarthritis of the knee: a 6 week double blind, placebo controlled pilot study of rofecoxib. *Rofecoxib Osteoarthritis Pilot Study Group*. *J Rheumatol* 1999;26:2438-2447.
176. Simon LS, Lanza FL, Lipsky PE, Hubbard RC. Preliminary study of the safety and efficacy of SC-58635, a novel cyclooxygenase 2 inhibitor: efficacy and safety in two placebo-controlled trials in osteoarthritis and rheumatoid arthritis and studies of gastrointestinal and platelet effects. *Arth Rheum* 1998;41:1591-1602.
177. Silverstein FE, Faich G, Goldstein JL et al. Gastrointestinal toxicity with celecoxib vs nonsteroidal anti-inflammatory drugs for osteoarthritis and rheumatoid arthritis: the CLASS study: A randomized controlled trial. *Celecoxib Long-term Arthritis Safety Study*. *JAMA* 2000;284:1247-1255.
178. Bombardier C, Laine L, A. R et al. Comparison of upper gastrointestinal toxicity of rofecoxib and naproxen in patients with rheumatoid arthritis. *N Engl J Med* 2000;343:1520-1528.
179. Simon LS. Toxicities of the nonsteroidal anti-inflammatory drugs. *Curr Opin Rheumatol* 1992;4:301-308.
180. Welch HG, Albertsen PC, Nease RF et al. Estimating treatment benefits for the elderly: the effects of competing risks. *Ann Intern Med* 1996;124:577-584.
181. Nicholas JJ. Physical modalities in rheumatological rehabilitation. *Arch Phys Med Rehabil* 1994;75:994-1001.
182. Deal CL, Schnitzer TJ, Lipstein E et al. Treatment of arthritis with topical capsaicin: a double-blind trial. *Clin Ther* 1991;13:383-395.
183. McCarthy GM, McCarty DJ. Effect of topical capsaicin in the therapy of painful osteoarthritis of the hands. *J Rheumatol* 1992;19:604-607.
184. Schnitzer TJ. Non-NSAID pharmacologic treatment options for the management of chronic pain. *Am J Med* 1998;105:45S-52S.
185. Kirwan J, Rankin E. Intra-articular therapy in osteoarthritis. *Bailliere's Clin Rheumatol* 1997;11:769-794.
186. Caldwell JR. Intra-articular corticosteroids: guide to selection and indications for use. *Drugs* 1996;52:507-514.
187. Adams ME, Atkinson MH, Lussier JS et al. The role of viscosupplementation with hylan G-F 20 (Synvisc) in the treatment of osteoarthritis of the knee: a Canadian multicenter trial comparing hylan G-F 20 alone, hylan G-F 20 with non-steroidal anti-inflammatory drugs (NSAIDs) and NSAIDs alone. *Osteoarth Cartilage* 1995;3:213-226.
188. Pozo MA, Balazs EA, Belmonte C. Reduction of sensory responses to passive movements of inflamed knee joints by hylan, a hyaluronan derivative. *Exp Brain Res* 1997;116:3-9.
189. Leardini G, Perbellini A, Franceschini M et al. Intra-articular injections of hyaluronic acid in the treatment of painful shoulder. *Clin Ther* 1988;10:521-526.
190. McAlindon TE, LaValley MP, Gulin JP et al. Glucosamine and chondroitin for treatment of osteoarthritis: a systematic quality assessment and meta-analysis. *JAMA* 2000;283:1469-1475.
191. daCamara CC, Dowless GV. Glucosamine sulfate for osteoarthritis. *Ann Pharmacother* 1998;32:580-587.